

# Multi-frequency emission analysis and constraining redshift of 3C 66A

Dahai Yan<sup>1\*</sup>, Zhonghui Fan<sup>1†</sup> and Benzhong Dai<sup>1‡</sup>

<sup>1</sup>*Department of Physics, Yunnan University, Kunming 650091, Yunnan, China*

Accepted –. Received –; in original form –

## ABSTRACT

The emission mechanism of TeV blazar 3C 66A is studied employing multi-wavelength observational data. The GeV spectrum observed by *Fermi*-LAT cannot be explained by pure synchrotron self-Compton (SSC) radiation. A multi-component leptonic jet model including the absorption of extragalactic background light reproduces the quasi-simultaneous spectral energy distribution. The external Compton component of the broad line region (BLR) is important for emission above 10 GeV, while emission from 0.1 GeV to 10 GeV is dominated by the SSC component. Our model implies that the redshift of 3C 66A should be between 0.15 and 0.31, the most likely value being 0.21. Our results also indicate that the  $\gamma$ -ray emission region is beyond the inner zone of BLR. The effects of model parameters and assumptions on the estimating of the redshift are discussed.

**Key words:** BL Lacertae objects: individual: 3C 66A – galaxies: active – gamma rays: general – radiation mechanisms: non-thermal.

## 1 INTRODUCTION

Blazars are the most extreme class of active galactic nuclei (AGN). Their spectral energy distributions (SEDs) are characterized by two distinct bumps. The low-energy component originates in a relativistic jet, and consists of relativistic electron synchrotron emission. The high-energy component could be produced by inverse Compton (IC) scattering (for a recent review, see Böttcher 2007). Various soft photon sources seed the SSC and EC processes in the jet to produce gamma rays.

Very high energy (VHE) photons emitted by extragalactic sources are absorbed, through the pair-production process, by interaction with extragalactic background light (EBL) (Stecker et al. 1992). The rate of absorption is determined by the EBL photon density and the redshift of the VHE source. The energy range of interest for background photons is from optical to ultraviolet (UV). Since it is hard to measure the EBL directly, researchers have proposed three kinds of spectral models: low limit (e.g., Kneiske et al. 2010; Razzaque et al. 2009), mean level (e.g., Finke et al. 2010a; Franceschini et al. 2008), and high level (e.g., Stecker et al. 2006). Aharonian et al. (2006) discussed some  $\gamma$ -ray blazars with unexpectedly hard spectra at large

redshifts, and suggested that EBL is of the first type. Albert et al. (2008) found that the universe is more transparent to  $\gamma$ -rays. However, Stecker et al. (2009) pointed out that Albert et al. (2008) do not significantly constrain the intergalactic low energy photon spectra and their high level EBL model is still valid. In an analysis of photons above 10 GeV from  $\gamma$ -ray sources detected by *Fermi*-LAT, Abdo et al. (2010a) found evidence to exclude the high level EBL model. By taking into account multi-frequency data ranging from radio to GeV, the SED of a blazar can be extrapolated into the VHE region. The redshift of the VHE source can then be constrained by comparing the de-absorbed VHE spectrum with the extrapolated one. The accurate determination of such redshifts helps us understand both the nature of EBL and the intrinsic emission model of VHE sources.

3C 66A is classified as intermediate BL Lac (IBL), because its synchrotron peak lies between  $10^{15}$  Hz and  $10^{16}$  Hz (Perri et al. 2003; Abdo et al. 2010b). The most widely used redshift for 3C 66A is 0.444, based on a single emission line measurement (Miller et al. 1978). However, Miller et al. (1978) stated that they were not sure of the reality of this emission feature, and warned that the redshift was not reliable. Later, Lanzetta (1993) confirmed the redshift of 0.444 based on data from *International Ultraviolet Explorer* (*IUE*). However, Bramel et al. (2005) argued that the 3C 66A redshift determined using *IUE* data is questionable. Finke et al. (2008a) placed a lower limit on the redshift of 3C

\* E-mail: yandahai555@gmail.com (DHY)

† E-mail: fanzh@ynu.edu.cn (ZHF)

‡ E-mail: bzhdai@ynu.edu.cn (BZD), Corresponding author

66A,  $z \geq 0.096$ , using information regarding its host galaxy. Recently, Prandini et al. (2010) concluded that the redshift of 3C 66A should be below  $0.34 \pm 0.05$ , and that the most likely redshift is  $0.21 \pm 0.05$ . These results assume that the de-absorbed VHE spectrum are not harder than the *Fermi*-LAT spectrum. Joshi & Böttcher (2007) argued that  $\gamma$ -ray emission in the flare state could be dominated by an external Compton (EC) process. Yang & Wang (2010) found that the VHE emission is contributed by EC when taking  $z = 0.444$ , or by pure SSC when  $z = 0.1$ . Note that this last result does not employ detailed *Fermi* data. Abdo et al. (2011) fit well the SED of 3C 66A at flare state with the SSC+EC model, and suggested that the redshift of 3C 66A may be between 0.2 and 0.3.

A quasi-simultaneous, multi-wavelength campaign of observations for 3C 66A was carried out by *Fermi* and *Swift* from August 2008 to November 2008. Data from the radio, optical, UV, X-ray, and GeV  $\gamma$ -ray to TeV  $\gamma$ -ray bands are available (Abdo et al. 2010b). VERITAS observed 3C 66A from September through November 2008 (Acciari et al. 2009a). Assuming the redshift of 3C 66A is 0.444, its VHE spectrum corrected using the mean level EBL model is harder than the observed GeV spectrum (Abdo et al. 2010b; Acciari et al. 2009a, 2010a). In this work, we fit the up-to-date SED of 3C 66A with a multi-component lepton jet model that takes into account EBL absorption. This model constrains the redshift to lie between 0.15 and 0.31. In Section 2, we describe the model and constrain its free parameters. Our results are presented in Section 3. A discussion of the data and our conclusions are given in Section 4. We adopt the cosmological parameters  $(H_0, \Omega_m, \Omega_\Lambda) = (70 \text{ km s}^{-1} \text{ Mpc}^{-1}, 0.3, 0.7)$  throughout.

## 2 MODEL AND PARAMETERS' CONSTRAINING

We assume that  $\gamma$ -ray emission is produced in a spherical blob of the jet, which is moving relativistically at a small angle to our line of sight. The observed radiation is strongly boosted by a relativistic Doppler factor  $\delta_D$ . The relativistic electrons inside the blob lose energy via synchrotron emission and IC scattering. The electron distribution is (Dermer et al. 2009),

$$N'_e(\gamma') = K'_e H(\gamma'; \gamma'_{\min}, \gamma'_{\max}) \gamma'^{-p_1} \exp(-\gamma'/\gamma'_b) \times H[(p_2 - p_1)\gamma'_b - \gamma'] + [(p_2 - p_1)\gamma'_b]^{p_2 - p_1} \gamma'^{-p_2} \times \exp(p_1 - p_2) H[\gamma' - (p_2 - p_1)\gamma'_b] K'_e H(\gamma'; \gamma'_{\min}, \gamma'_{\max}), \quad (1)$$

where  $H(x; x_1, x_2)$  is the Heaviside function:  $H(x; x_1, x_2) = 1$  for  $x_1 \leq x \leq x_2$  and  $H(x; x_1, x_2) = 0$  everywhere else; as well as  $H(x) = 0$  for  $x < 0$  and  $H(x) = 1$  for  $x \geq 0$ . In the co-moving frame, this distribution is a double power law with two energy cutoffs:  $\gamma'_{\min}$  and  $\gamma'_{\max}$ . The spectrum is smoothly connected, with indices  $p_1$  and  $p_2$  below and above the electrons' break energy  $\gamma'_b$ . Note that here and throughout the paper, unprimed quantities refer to the observer's frame and primed ones refer to the co-moving frame.

The multi-component model of Dermer et al. (2009) is used to calculate the multi-wavelength SED. This model takes into account  $\gamma$ -ray attenuation by the BLR-scattered radiation field. For EC components, we con-

sider photons emitted directly from the accretion disk, using BLR-scattered photons as the seed photons. Here we briefly describe the model for numerical calculation, please see Dermer et al. (2009) in detail. The  $\nu F_\nu$  spectrum of Compton-scattered external photons is

$$f_{\epsilon_s}^{\text{EC}} = \frac{c}{4\pi d_L^2} \epsilon_s^2 \delta_D^3 \int_0^{2\pi} d\phi_* \int_{-1}^1 d\mu_* (1 - \mu_*) \times \int_0^{\epsilon_{*,\text{up}}} d\epsilon_* \frac{u_*(\epsilon_*, \Omega_*)}{\epsilon_*} \int_{\gamma_{\text{low}}}^\infty d\gamma N'_e(\gamma/\delta_D) \frac{d\sigma_C}{d\epsilon_s}, \quad (2)$$

where  $\gamma = \gamma' \delta_D$ ,  $d_L(z)$  is the luminosity distance,  $z$  is the source redshift,  $c$  is the speed of light,  $\mu_*$  is the collision angle between the relativistic electrons and seed photons and  $u(\epsilon_*, \Omega_*)$  is energy density of external photons with energy  $\epsilon_*$ . The Compton cross section in the head-on approximation, accurate for relativistic electrons from the Thomson through the Klein-Nishina regime, is

$$\frac{d\sigma_C}{d\epsilon_s} \cong \frac{\pi r_e^2}{\gamma \bar{\epsilon}} [y + y^{-1} - \frac{2\epsilon_s}{\gamma \bar{\epsilon} y} + \frac{\epsilon_s^2}{\gamma^2 \bar{\epsilon}^2 y^2}] H(\epsilon_s; \frac{\bar{\epsilon}}{2\gamma}, \frac{2\gamma \bar{\epsilon}}{1 + 2\bar{\epsilon}}) \quad (3)$$

where  $r_e$  is the classical electron radius, the invariant collision energy  $\bar{\epsilon} = \gamma \epsilon_* (1 - \mu_*)$  and  $y = 1 - \epsilon_s/\gamma$ . The lower limit on the electron energy  $\gamma_{\text{low}}$  and the upper limit  $\epsilon_{*,\text{up}}$  are

$$\gamma_{\text{low}} = \frac{\epsilon_s}{2} [1 + \sqrt{1 + \frac{2}{\epsilon_* \epsilon_s (1 - \mu_*)}}], \quad (4)$$

and

$$\epsilon_{*,\text{up}} = \frac{2\epsilon_s}{1 - \mu_*}. \quad (5)$$

Gamma-ray photons are subject to  $\gamma\gamma$  attenuation by the ambient photons in the environment of the black hole. The absorption probability per unit pathlength is

$$\frac{d\tau^{\text{internal}}(\epsilon_1)}{dl} = \oint d\Omega_* (1 - \mu_*) \int_0^\infty \frac{u(\epsilon_*, \Omega_*)}{m_e c^2 \epsilon_*} \sigma_{\gamma\gamma}(\beta) d\epsilon_* \quad (6)$$

(Dermer et al. 2009), where  $\mu_*$  is the angle between the direction of the Gamma-ray photon with energy  $\epsilon_1$  and the target photon with energy  $\epsilon_*$  and  $m_e$  is the rest mass of electron (the scattered photon travels in the same direction as the relativistic electron). The cross-section  $\sigma_{\gamma\gamma}(\beta)$  is given by Gould & Shréder (1967), with  $\beta = \sqrt{1 - 1/s}$  and  $s = \epsilon_1 \epsilon_* (1 + z)(1 - \mu_*)/2$ .

We assume that the BLR is a spherically symmetric shell with inner radius  $R_i$  and outer radius  $R_o$ . We also assume that the gas density of the BLR has the power-law distribution  $n_e(r) = n_0 (\frac{r}{R_i})^\zeta$ , where  $R_i \leq r \leq R_o$ . The radial Thomson depth is given by  $\tau_T = \sigma_T \int_{R_i}^{R_o} dr n_e(r)$ , where  $r$  is the distance from the central black hole (Dermer et al. 2009). In our calculation we use  $\tau_T = 0.01$ , which is the typical value (e.g., Finke et al. 2010b; Reimer 2007; Donea & Protheroe 2003). Kaspi & Netzer (1999) suggested that the particle density of BLR scales as  $r^{-1.0}$  or  $r^{-1.5}$ . In our calculation, we adopt the exponent  $\zeta = -1.0$ .

Using reverberation mapping, Bentz et al. (2009) derived an improved empirical relationship between BLR radius  $R_{\text{BLR}}$  and luminosity  $L_\lambda$  at 5100Å:

$$\log(R_{\text{BLR}}) = -21.3 + 0.519 \cdot \log(\lambda L_\lambda(5100\text{\AA})). \quad (7)$$

The V-band magnitude of 3C 66A is 15.21

(Véron-Cetty & Véron 2010). We use the optical spectral index given by Fiorucci et al. (2004) to calculate the average flux at 5100Å, which is 2.785 mJ. In this work, we take the estimated  $R_{\text{BLR}}$  as the outer radius of the BLR  $R_o$ . Peterson et al. (1994) suggested that the typical size of the BLR in quasars is on the order of light-months. We follow several authors (e.g., Reimer 2007; Donea & Protheroe 2003), using the relationship  $R_i = R_o/40$  to derive a value for  $R_i$ .

To simplify calculation, the BLR-scattered photon field is assumed to be monochromatic with energy  $\epsilon_*$ , which is the mean energy from the accretion disk (Dermer et al. 2009). The approximate mean dimensionless photon energy at radius  $R$  from a standard accretion disk is given by (Finke et al. 2010b)

$$\epsilon_d(R) = 1.5 \times 10^{-4} \left( \frac{10\ell_{\text{Edd}}}{M_8\eta} \right)^{\frac{1}{4}} \left( \frac{R}{r_g} \right)^{-\frac{3}{4}}. \quad (8)$$

The accretion luminosity is  $\ell_{\text{Edd}} = \frac{L_d}{L_{\text{Edd}}}$ , which here has the value 0.03. The Eddington luminosity is  $L_{\text{Edd}} = 1.26 \times 10^{46} M_8 \text{ ergs} \cdot \text{s}^{-1}$ , and  $L_d$  is the accretion disk luminosity. The accretion efficiency  $\eta$  is 0.1. The gravitational radius is  $r_g = \frac{GM}{c^2} \cong 1.5 \times 10^{13} M_8 \text{ cm}$ . The mass of the 3C 66A black hole is  $M_8 = \frac{M_{\text{BH}}}{10^8 M_\odot} = 4.0$  (Ghisellini et al. 2010). The typical energy of photons from the disk is  $\epsilon_d(2.25R_{\text{in}}^d)$  (Shakura & Sunyaev 1973), where  $R_{\text{in}}^d$  is the inner radius of accretion disk.  $R_{\text{in}}^d$  is approximately equal to  $3r_g$  for a Kerr black hole, or  $6r_g$  for a Schwarzschild black hole. In this work, we adopt  $\epsilon_* = \epsilon_d(10r_g) = 2.48 \times 10^{-5}$ . The energy density of BLR-scattered photon field is

$$u(\epsilon_*, \mu_*; r_b) = \frac{L_d r_e^2}{3ct_b} F(\mu_*, r_b) \quad (9)$$

(Dermer et al. 2009), where  $r_b$  is the distance from the emission blob to the central black hole and  $F(\mu_*, r_b)$  is the function given by Dermer et al. (2009) (their Eq.(97)).

The intrinsic flux from extragalactic sources is

$$f_{\text{intrinsic}}(E_\gamma) = e^{\tau(E_\gamma, z)} f_{\text{observed}}(E_\gamma), \quad (10)$$

where  $f_{\text{observed}}$  is the measured VHE flux, and  $\tau(E_\gamma, z)$  is the optical depth of a  $\gamma$ -ray with energy  $E_\gamma$  at redshift  $z$ .  $\tau(E_\gamma, z)$  is given by (Franceschini et al. 2008)

$$\tau(E_\gamma, z) = c \int_0^z dz_1 \frac{dt}{dz_1} \int_0^2 dx \frac{x}{2} \times \int_{2m_e^2 c^4 / E_\gamma (1+z_1)x}^\infty d\epsilon \frac{dn(\epsilon, z_1)}{d\epsilon} \sigma_{\gamma\gamma}(\beta). \quad (11)$$

The derivative  $\frac{dt}{dz_1}$  gives the relationship between cosmic time and redshift, which is  $\frac{dt}{dz_1} = \frac{1}{H_0(1+z_1)} [(1+z_1)^2 (\Omega_m z_1 + 1) - z_1(z_1+2)\Omega_\Lambda]^{-1/2}$ .  $\frac{dn(\epsilon, z_1)}{d\epsilon}$  is the number density of EBL photons with energy  $\epsilon$  at redshift  $z_1$ , and  $x = 1 - \cos\theta$ , where  $\theta$  is the collision angle.

Here, we use the EBL model of Franceschini et al. (2008)<sup>1</sup> to de-absorb the observed VHE SED. This model is based on observations and takes into account all available

**Table 1.** Model Parameters for Fig. 1.

| parameters                         | $z = 0.15$ | $z = 0.21$ | $z = 0.31$ |
|------------------------------------|------------|------------|------------|
| $\gamma'_b (10^3)$                 | 5.8        | 6.3        | 7.6        |
| $\gamma'_{\text{min}} (10^3)$      | 1.93       | 1.90       | 1.76       |
| $\delta_D$                         | 38         | 36         | 43         |
| $t_{v,\text{min}} (10^4 \text{s})$ | 0.69       | 1.17       | 1.21       |
| $R_o$ (pc)                         | 0.1        | 0.14       | 0.22       |
| $r_b$ ( $R_o$ )                    | 1.03       | 0.89       | 0.72       |
| $B/B_{\text{eq}}$                  | 0.306      | 0.335      | 0.352      |
| $\chi^2$                           | 3.099      | 1.498      | 3.880      |

information on cosmic sources contributing background photons. Franceschini et al. (2008) also considered the redshift evolution of EBL.

Several parameters in our model can be constrained by observations. Böttcher et al. (2005) estimated the Doppler factor of 3C 66A as  $\delta_D \approx 15$ . Böttcher et al. (2009) excluded extreme values of the Doppler factor in the range  $\delta_D \geq 50$ . The size of the emission blob can be constrained by the observed variability timescales  $t_{\text{var}}$ , because  $R_b = t_{v,\text{min}} \delta_D c / (1+z) \leq \delta_D c t_{\text{tar}} / (1+z)$ . Here  $R_b$  is the radius of the blob in the comoving frame, and  $t_{v,\text{min}}$  is the smallest variability timescale. Takalo (1996) reported a micro-variability with  $t_{\text{var}} \sim 2.16 \times 10^4 \text{s}$  and  $\Delta \text{mag} \sim 0.2$ . Abdo et al. (2011) reported shorter variability at optical band:  $t_{\text{var}} \sim 1.44 \times 10^4 \text{s}$ .

### 3 THE RESULTS

From the observed photon index in the X-ray band  $\Gamma_X$  (Abdo et al. 2010b), we can roughly estimate that  $p_2 \approx 4.0$  using the relationship  $\Gamma_X - 1 = \frac{1}{2}(p_2 - 1)$ . In our calculation, the values of  $p_1$  and  $p_2$  are fixed at 2.0 and 4.0 respectively. There are few constraints on the values of the magnetic field  $B$  and  $\gamma_{\text{max}}$ . We use the fixed parameters  $B = 0.168 \text{ G}$  and  $\gamma_{\text{max}} = 3.0 \times 10^6$ . Table 1 lists all other model parameters, including  $B_{\text{eq}}$  (the equipartition magnetic field). The reduced  $\chi^2$  is calculated from the normalized residuals.

In Fig. 1, we show the modeling results at different plausible redshifts. The filled circles are quasi-simultaneous data in bands ranging from radio to GeV. The observed VERITAS data are de-absorbed using the EBL model of Franceschini et al. (2008) with different redshifts. We obtain the best fit when  $z = 0.21$ . It is found that the accretion-disk component is negligible compared to the SSC and BLR components. SSC and EC are responsible for emission in the GeV-TeV range. Emission from 0.1 GeV to 10 GeV is dominated by SSC. Above 10 GeV, the EC component of BLR is more important.

In Fig. 2, we show the  $\gamma\gamma$  absorption by BLR-scattered radiation at different locations of the blob when taking  $z = 0.21$ . There is significant absorption when the blob is inside the inner zone of the BLR. Beyond the inner zone, absorption is negligible.

<sup>1</sup> Opacities for photon-photon interaction as a function of the source redshift are available on the the website <http://www.astro.unipd.it/background>.




**Figure 1.** In panel (a), we show the best modeling results with  $z = 0.21$ . The filled square are the de-absorbed VHE data with  $z = 0.21$ . The dashed, dash-dotted, dotted and thick solid lines are SSC component, accretion-disk, BLR-reproduced component and the sum of multi-component, respectively. In panel (b), the open square, filled square and open circle are the de-absorbed VHE data with  $z = 0.15, 0.21$  and  $0.31$ , respectively. The dash-dotted, solid and dashed lines are the model results at  $z = 0.15, 0.21$  and  $0.31$ , respectively. All observed data are from Abdo et al. (2010b).




**Figure 2.**  $\gamma\gamma$  optical depth for  $\gamma$ -ray interaction with BLR-reproduced photons at different distances from central BH when  $z = 0.21$ .

#### 4 DISCUSSION AND CONCLUSIONS

At present, three IBLs (3C 66A, W com and PKS 1424+240) have been detected in VHE region. The first two require an EC component to model high-energy emission (e.g., Acciari et al. 2009b, 2010b; Yang & Wang 2010). A pure SSC model fails to explain the GeV spectrum of 3C 66A observed by *Fermi*-LAT. However, a satisfactory fit to these data can be obtained by our multi-component model (see Fig. 1).

The multi-component model requires a large value of  $\gamma'_{\min} \sim 2 \times 10^3$ . As argued by Tavecchio et al. (2009), this outcome seems to provide important clues to the electron

acceleration process and the role of energy loss. A large value of  $\gamma'_{\min}$  leads to a steep spectrum in the low-energy band, so our model does not explain the observed radio emission. This may come from a larger emission region. Assuming the total particle power is ten times of the electron power in the jet (Finke et al. 2008b), the ratio between the best-fit magnetic field and equipartition magnetic field (see Table 1) is close to a state of equipartition between the total particle energy density and the magnetic field energy density.

A simple assumption is that the BLR scattered photon field is characterized by an isotropic blackbody (e.g., Inoue & Takahara 1996; Yang & Wang 2010). Our model takes into account not only the specific shell structure of the BLR, but also the angular dependence of the photon distribution. As shown in panel (a) of Fig. 1, BLR-scattered photons are important to the EC process. Fan et al. (2006) also found that soft seed photons for EC predominantly come from the BLR in a sample of  $\gamma$ -ray loud blazars. Furthermore, our results show that the contributions of different EC components to the  $\gamma$ -ray spectrum are dependent on the location of the blob. As the distance between the central BH and the blob increases, the BLR component grows more important relative to the accretion-disk component. The BLR-scattered radiation field not only provides the seed photons for IC, but also attenuates the  $\gamma$ -rays. In flat-spectrum radio quasars (FSRQs), absorption by BLR radiation is significant and can cause a break in the GeV band (e.g., Reimer 2007; Sitarek & Bednarek 2008; Bai et al. 2009; Poutanen et al. 2010). Our results show that  $\gamma$ -ray absorption by BLR radiation becomes more and more slight as the value of  $r_b$  increases (see Fig. 2).

Thanks to these high-quality, multi-frequency data, we can use the extrapolated intrinsic VHE spectra to constrain the redshift of 3C 66A. As shown in panel (b) of Fig. 1, the de-absorbed VHE spectrum is too steep to be fitted well by our model if the redshift is below 0.15. However, the same is true if the redshift is above 0.31, as the shape of the de-absorbed VHE spectra becomes strange. Hence, the redshift of 3C 66A should be between 0.15 and 0.31. We calculate the reduced  $\chi^2$  for 11  $\gamma$ -ray spectra at different redshifts between 0.15 and 0.31. The minimum value,  $\chi^2 = 1.498$ , occurs for  $z = 0.21$ , which means the most likely redshift for 3C 66A is 0.21. Prandini et al. (2010) assume that the VHE spectrum corrected for the absorption through the interaction with the EBL can not be harder than the spectrum in the Fermi/LAT band. They derived a EBL-corrected spectra as a function of the redshift with a TeV blazars sample of known redshift. Comparing the TeV corrected spectra with GeV LAT spectra, they get the estimation distance of unknown redshift blazars. They suggested that the most likely redshift of 3C 66A is  $0.21 \pm 0.05$ . Two results from different methods are completely consistent.

The contribution of the BLR component is crucial for constraining the redshift of 3C 66A. The BLR structure ( $R_i$ ,  $R_o$ ,  $\zeta$ ,  $\tau_T$ ) and the characteristics of the central source (the black hole and its accretion disk) can affect the contribution of the BLR component.  $R_o$  can be constrained by Eq.(7). We assume  $(\ell_{\text{Edd}}, R_o/R_i, \tau_T, \zeta) = (0.03, 40, 0.01, -1)$  and get the best model fitting with  $z = 0.21$ . The effects of these parameters and assumptions on the estimating of the redshift are discussed. Fig. 3(a), (b), (c) and (d) show the effects of accretion luminosity, BLR size, radial Thomson depth and

**Table 2.** Model Parameters for Fig. 3. The blank ones represent these parameter are the same as that we adopted when  $z = 0.21$ .

| parameters                       | $\ell_{\text{Edd}} = 0.01$ | $\frac{R_o}{R_i} = 5$ | $\tau_T = 0.1$ | $\zeta = -2$ |
|----------------------------------|----------------------------|-----------------------|----------------|--------------|
| $\gamma'_b(10^3)$                |                            |                       | 5.6            |              |
| $\gamma'_{\text{min}}(10^3)$     | 1.8                        | 2.0                   | 2.5            |              |
| $\delta_D$                       |                            |                       | 37             |              |
| $t_{v,\text{min}}(10^4\text{s})$ | 1.2                        |                       | 1.05           | 3.2          |
| $r_b(R_o)$                       | 0.65                       | 1.02                  | 1.31           | 0.52         |
| $B/B_{\text{eq}}$                | 0.333                      | 0.344                 | 0.366          | 0.794        |
| $\chi^2$                         | 1.251                      | 1.511                 | 1.934          | 1.314        |

distribution exponent of BLR gas density on the model fitting respectively. For clarity, only the modeling results in the high-frequency range of the case of  $z = 0.21$  are shown. The model parameters are listed in Table 2. The parameters which we don't particularly claim are the same as that we used in the case of  $z = 0.21$ . The  $\gamma\gamma$  absorption by BLR-scattered radiation is also negligible in all of the four cases.

From the reduced  $\chi^2$  (see Table 2), we find the fitting results are in agreement with that we originally obtained. We therefore conclude that the assumptions of BLR structure and the characteristics of the central source don't affect our estimating on the redshift. It can be interpreted as that these assumptions affect the density of external photons field, and we can adjust the location of the blob ( $r_b$ ) to determine the suitable density of external photons field to reproduce the GeV radiation. Therefore, determining the location of the blob is important for constraining the structure of the BLR. It's found that the reduced  $\chi^2$  gets better when  $\ell_{\text{Edd}} = 0.01$  and worse when  $\tau_T = 0.1$  compared with the case of  $\ell_{\text{Edd}} = 0.03$  and  $\tau_T = 0.01$ , respectively. Hence, the more appropriate parameters may be  $\ell_{\text{Edd}} = 0.01$  and  $\tau_T = 0.01$  for 3C 66A. From Table 2, we see that the changes of parameters are slight and reasonable except the case when  $\zeta = -2$  (see Table. 2). In the case of  $\zeta = -2$ , the density of the BLR-scattered photon field declines more quickly with the blob's distance from the central black hole. For fitting the GeV data, the smaller value of  $r_b$  is required, then the accretion-disk component becomes more important. As shown in panel (d) in Fig. 3, radiation from 0.1-10 GeV is dominated by accretion-disk component. In this case, larger value of  $t_{v,\text{min}} = 3.2 \times 10^4$  s is required, which is in contrast with the observed optical variability time scale:  $t_{\text{var}} \sim 1.44 \times 10^4$  s. Our results therefore indicate that the case of  $\zeta = -2$  seems to be excluded for 3C 66A. We also conclude from model fitting that the blob is located beyond the inner zone of the BLR, may be between  $0.5R_o$  and  $1.3R_o$ .

In conclusion, we study the multi-frequency emission of 3C 66A by using multi-wavelength observational data over radio through GeV-TeV energy region. Under the mean-limit EBL absorption, the model constrains the redshift of 3C 66A to lie between 0.15 and 0.31. The most likely value is 0.21.




**Figure 3.** The effects of different assumptions of BLR structure and the characteristics of central source on the estimating of the redshift. The symbols are the same as that in Fig. 1.

## ACKNOWLEDGMENTS

We thank X. W. Cao for helpful comments on this paper and J. P. Yang for helpful discussion. This work is supported by the National Science Foundation of China (grants 11063003 and 10963004), Yunnan Provincial Science Foundation of China (grant 2008CD061) and SRFDP of China (grant 20095301120006).

## REFERENCES

- Abdo A. A. et al., 2010a, *ApJ*, 723, 1082
- Abdo A. A. et al., 2010b, *ApJ*, 716, 30
- Abdo A. A. et al., 2011, *ApJ*, 726, 43
- Acciari V. A. et al., 2009a, *ApJ*, 693, L104
- Acciari V. A. et al., 2009b, *ApJ*, 707, 612
- Acciari V. A. et al., 2010a, *ApJ*, 721, L203 (Erratum to *ApJ*, 693, L104)
- Acciari V. A. et al., 2010b, *ApJ*, 708, 100
- Aharonian F. et al., 2006, *Nat*, 440, 1018
- Albert J. et al., 2008, *Sci*, 320, 1752
- Bai J. M., Liu H. T., & Ma L., 2009, *ApJ*, 699, 2002
- Bentz M. c. et al., 2009, *ApJ*, 697, 160
- Böttcher M., 2007, *Ap&SS*, 309, 95
- Böttcher M. et al., 2005, *ApJ*, 631, 169
- Böttcher M. et al., 2009, *ApJ*, 694, 174
- Bramel D. A. et al., 2005, *ApJ*, 629, 108
- Dermer C. D., Finke J. D., Krug H., & Böttcher M., 2009, *ApJ*, 692, 32
- Donea A. C., & Protheroe R. J., 2003, *APh*, 18, 377
- Fan Z. H., Cao X. W., & Gu M. F., 2006, *ApJ*, 646, 8
- Finke J. D., Shields J. C., Böttcher M., & Basu S., 2008a, *A&A*, 477, 513
- Finke J. D., Dermer C. D., & Böttcher M., 2008b, *ApJ*, 686, 181
- Finke J. D., Razzaque S., & Dermer C. D., 2010a, *ApJ*, 712, 238
- Finke J. D., & Dermer C. D., 2010b, *ApJ*, 714, 303

- Fiorucci M., Ciprini S., & Tosti G., 2004, *A&A*, 419, 25
- Franceschini A., Rodighiero G., & Vaccari M., 2008, *A&A*, 487, 837
- Ghisellini G. et al., 2010, *MNRAS*, 402, 497
- Gould R. J., & Shröder G. P., 1967, *Phys. Rev.*, 155, 1404
- Inoue S., & Takahara F., 1996, *ApJ*, 463, 555
- Joshi M., & Böttcher M., 2007, *ApJ*, 662, 884
- Kaspi S., & Netzer H., 1999, *ApJ*, 524, 71
- Kneiske T. M., & Dole H., 2010, *A&A*, 515, 19
- Lanzetta K. M., Turnshek D. A., & Sandoval J., 1993, *ApJS*, 84, 109
- Miller J. S., French H. B., & Hawley S. A., 1978, in *Pittsburgh Conference on BL Lac Objects*, ed. A. M. Wolfe (Pittsburgh, PA: Univ. Pittsburgh), 176
- Perri M. et al., 2003, *A&A*, 407, 453
- Peterson B. M. et al., 1994, *ApJ*, 425, 622
- Poutanen J., & Stern B., 2010, *ApJ*, 717, 118
- Prandini E., Bonnoli G., Maraschi L., Mariotti M., & Tavecchio F., 2010, *MNRAS*, 405, L76
- Razzaque S., Dermer C. D., & Finke J. D., 2009, *ApJ*, 697, 483
- Reimer A., 2007, *ApJ*, 665, 1023
- Shakura N. I., & Sunyaev R. A., 1973, *A&A*, 24, 337
- Sitarek J., & Bednarek W., 2008, *MNRAS*, 391, 624
- Stecker F. W., de Jager O. C., & Salamon M. H., 1992, *ApJ*, 390, L49
- Stecker F. W., Malkan M. A., & Scully S. T., 2006, *ApJ*, 648, 774
- Stecker F. W., & Scully S. T., 2006, *ApJ*, 691, L91
- Takalo L. O. et al., 1996, *A&AS*, 120, 313
- Tavecchio F., Ghisellini G., Ghirlanda G., Costamante L., & Franceschini A., 2009, *MNRAS*, 399, 59
- Véron-Cetty M. P., & Véron P., 2010, *A&A*, 518A, 10
- Yang J. P., & Wang J. C., 2010, *A&A*, 511, 11

This paper has been typeset from a  $\mathrm{T}_{\mathrm{E}}\mathrm{X}/\mathrm{L}^{\mathrm{A}}\mathrm{T}_{\mathrm{E}}\mathrm{X}$  file prepared by the author.